

Diaper Rash

Definition

- Any rash on the skin covered by a diaper
- Diaper-wearing age group (birth to 3 years)

Health Information

Symptoms

- Mild rashes just have areas of pink dry skin.
- Severe rashes have areas of red skin. In some areas, the skin may become raw or even bleed.
- Yeast infections are bright red. The borders are very sharp. Small red bumps or even pimples may occur just beyond the border. Yeast rashes usually cover large areas.
- Pink rashes are not painful, but raw ones can be very painful. This can lead crying and poor sleep.

Cause

- Mild rashes can be caused by the drying effect of soaps.
- Stool and urine left on skin can combine to make ammonia. Ammonia can cause a mild chemical burn.
- Stools left on the skin can be very irritating because they contain bacteria. Urine alone has no germs in it and usually doesn't irritate the skin.
- Rashes around the anus are common during bouts of diarrhea.
- Rashes from irritants can get a secondary infection with yeast. Less commonly it can happen with bacteria.

Prevention of Recurrent Diaper Rash

- Change diapers more often. Especially try to prevent skin contact with stool.
- Rinse the baby's skin with lots of warm water when cleaning off stool. Don't depend on diaper wipes alone
- Be sure to clean stool off all the skin folds. Cleaning the scrotum can be a challenge.

Care Advice

1. Overview:

- Diaper rashes are very common in babies.
- Often caused by not cleaning stool off the skin soon enough.
- Here's some care advice that should help.

2. Change More Often:

- Change diapers more often to prevent skin contact with stool.
- You may want to get up once during the night to change the diaper. Do this until the rash is under control.

3. Rinse with Warm Water:

- Rinse the baby's skin with lots of warm water after cleaning off stool.
- Wash with a mild soap (such as Dove) only after stools. Reason: Using soap often can slow healing.
- Do not depend on diaper wipes alone for removing stool. Reason: They leave a film of bacteria on the skin.

4. **Leave the Bottom Open to Air:**
 - Expose the bottom to air as much as possible.
 - Attach the diaper loosely at the waist to help with air exposure.
 - When napping, take the diaper off and lay your child on a towel. Reason: Dryness reduces the risk of yeast infections.
5. **Anti-Yeast Cream:**
 - Most diaper rashes respond to 3 days of warm water cleansing and air exposure. If you've tried this or the rash is bright red, suspect a yeast infection.
 - Buy an anti-yeast cream such as Lotrimin. No prescription is needed.
 - Use Lotrimin cream 2 times per day.
6. **Raw Skin:**
 - If the bottom is very raw, soak in warm water for 10 minutes. Add 2 tablespoons (30 ml) of baking soda to the tub of warm water.
 - Do this 2 times per day for a few days.
 - Then, put Lotrimin cream on the rash.
7. **Sore or Scab on End of the Penis:**
 - Use an antibiotic ointment such as Polysporin. No prescription is needed.
 - Do this 3 times per day.
 - Reason: The sore is a bacterial infection that can cause painful urination.
8. **Diarrhea Rash:**
 - If your child has diarrhea and an anal rash, use a protective ointment. Examples are petroleum jelly, A&D or Desitin.
 - This forms a barrier between the skin and the stool.
 - Otherwise, these generally are not needed.
 - Caution: Wash off the skin before putting the ointment on.
9. **What to Expect:**
 - With proper treatment, most diaper rashes are better in 3 days.
 - If the rash does not respond, a yeast infection has probably occurred. If so, start an anti-yeast cream.
10. **Return to Child Care:**
 - Diaper rashes cannot be spread to others. Yeast ones do not spread. Bacterial ones have a small risk until they are on an antibiotic ointment.
 - Your child does not need to miss any child care.

Call Your Doctor If

- Rash isn't much better after 3 days of using anti-yeast cream
- It starts to look infected (with sores and scabs)
- You think your child needs to be seen
- Your child becomes worse

Pediatric Care Advice

Author: Barton Schmitt MD, FAAP

Copyright 2000-2022 Schmitt Pediatric Guidelines LLC

Disclaimer: This health information is for educational purposes only. You the reader assume full responsibility for how you choose to use it. The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. Listing of any resources does not imply an endorsement.